

# Adult-onset Still's disease and dermatomyositis, is there a relation?

Doaa Atef, Asmaa Sabbah, Shaimaa I. El-Jaafary, Noha T. AboKrysha

Department of Neurology, Faculty of Medicine, Cairo University, Cairo, Egypt

Correspondence to Asmaa Sabbah, MD, 829, District 1, Nighborhood 4, 6th October city, 12591, Egypt; Tel: +20 100 376 0790; e-mail: doctor\_sabbah@cu.edu.eg

Received 18 June 2016

Accepted 24 October 2016

Kasr Al Ainy Medical Journal  
2017, 23:63–65

Adult-onset Still's disease (AOSD) is a rare systemic inflammatory disorder of unknown etiology. We report a 22-year-old female patient, admitted to our department with fever, rash, large joint pains, and mild proximal weakness of both upper and lower limbs. With further evaluation of the patient and exclusion of infections, hematological malignancies, and connective tissue disorders, AOSD diagnosis was established with strikingly elevated serum ferritin level. This patient is the daughter of a patient who was diagnosed as having dermatomyositis in our department 2 years earlier. The mother was re-evaluated, and AOSD could not be excluded through the normal serum ferritin level.

## Keywords:

adult-onset Still's disease, dermatomyositis, myopathy, neurological manifestations

Kasr Al Ainy Med J 23:63–65  
© 2017 Kasr Al Ainy Medical Journal  
1687-4625

## Introduction

Adult-onset Still's disease (AOSD) is a rare systemic inflammatory disorder of unknown etiology that is frequently underdiagnosed and typically characterized by a clinical triad of daily spiking high fever, evanescent rash (salmon colored), and arthritis. A biological triad of hyperferritinemia, hyperleukocytosis with neutrophilia, and abnormal liver function test result also characterizes the disease.

There is no specific test or a combination of tests that establishes the diagnosis of AOSD; however, the clinical presentation together with elevated serum ferritin is highly suggestive of AOSD [1].

The incidence of AOSD is estimated to be 0.16 per 100 000 persons, with a bimodal peak at ages 15–25 and 36–46 years (with three-quarters of the patients reporting disease onset between 16 and 35 years of age) and equal distribution between males and females [2].

Although the exact etiology is still unknown, some observations supported a genetic predisposition and environmental factors that trigger the immune system (immune mediated disease); a range of infectious agents may trigger the disease in a genetically predisposed host [1].

## Case report

A 22-year-old female patient presented with multiple large joint pains and proximal weakness of both upper and lower limbs. She had spiking fever as well, especially in the evening reaching 39.5°C, associated with severe sore throat 2 weeks before the onset. The

onset of fever was also associated with an evanescent, pruritic erythematous linear rash mainly over the trunk and extremities. Her medical history was unremarkable, and her mother was a known case of dermatomyositis.

Examination revealed fever of 39.0°C. There was lymphadenopathy but no splenomegaly. She had acute synovitis of wrists, elbows, ankles, and knees. There was a residual skin rash seen over the trunk and extremities.

Neurological examination finding was free of any abnormalities apart from muscle tenderness and mild proximal weakness in both upper and lower limbs grade 4, with brisk deep tendon reflexes.

Hematological investigations showed leukocytosis of  $8 \times 10^9/l$  with neutrophilia. Erythrocyte sedimentation rate was high 80/115 mm/h. There were markedly elevated levels of serum ferritin (1616.6 ng/ml). The findings of antinuclear antibody, antidouble-strand DNA, and rheumatoid factor were all negative. Renal and liver function findings as well as coagulation profiles were normal. Blood and urine cultures revealed no evidence of bacterial, fungal, or viral infection.

Electromyogram showed evidence of myopathy.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work noncommercially, as long as the author is credited and the new creations are licensed under the identical terms.

Based on the clinical features and laboratory results, she was diagnosed as having AOSD according to the Yamaguchi criteria [3].

The patient was treated with corticosteroids, initially intravenously and then orally, with improvement in her symptoms.

The mother of our patient who was diagnosed as having dermatomyositis 2 years ago, with typical skin eruption and inflammatory myopathy, was asked to visit us to undergo laboratory tests, serum ferritin together with erythrocyte sedimentation rate, antinuclear antibody, and rheumatoid factor. The results of which came out negative. We revised the literature aiming at finding relation between AOSD and dermatomyositis, and there were some reports of AOSD presented with dermatomyositis like skin rash [4].

Having low serum ferritin level does not exclude the diagnosis of the mother as having AOSD as ferritin level is lower in remission, and in ~10% of the patients with AOSD, the ferritin levels are within reference range [5].

Although it is known that AOSD is a sporadic noninherited disease, we report a patient and her mother who had myopathic syndromes with skin rash; for the patient, the most likely diagnosis is AOSD.

---

## Discussion

AOSD is a rare inflammatory disorder that has been recently classified as a polygenic autoinflammatory disorder. AOSD can be divided into two phenotypes based on cytokine profile, 'systemic' and an 'articular' pattern. Interleukin-1 (IL-1) family (namely, IL-1 $\beta$  and IL-18) plays a major role in the pathophysiology of the systemic pattern of the disease. Drugs that antagonize IL-1 $\beta$ , like anakinra, have been used successfully in the treatment of resistant cases of AOSD [6]. Revising the literature revealed that IL-1 $\beta$  expression is one of the most consistent cytokine patterns in the muscle tissue of patients with dermatomyositis. Treatment with anakinra had a beneficial effect in a series of patients with refractory DM [7]. So it is clear now that both diseases share common pathophysiological aspects.

AOSD has different neurological presentations. The nervous system was affected in 7–12% of cases, mainly as cranial nerve palsies, pyramidal and extrapyramidal

affection, aseptic meningoencephalitis, demyelinating encephalopathy, seizures, myopathy, and Miller-Fisher syndrome have been also described [8–10].

The diagnosis is often difficult as the condition is similar to systemic infections, vasculitis, or even some malignant condition; to establish the diagnosis of AOSD, these condition should be excluded, and elevated levels of serum ferritin may help [3]. Our patient presented with almost all the major and minor criteria together with abnormally high serum ferritin level, and her mother has been diagnosed and treated as diabetes mellitus.

---

## Conclusion

AOSD can present for the first time at the neurology clinic and should be included in the differential diagnosis of young patients presenting with high fever with muscle pain and weakness associated with arthritis. The diagnosis of AOSD is done by exclusion after investigations for other similar conditions, for example, infections, granulomatous disorders, connective tissue disorders, and malignancies like lymphoma.

## Acknowledgements

The article has been read and approved by all the authors, the requirements for authorship as stated earlier in this document have been met, and each author believes that the article represents honest work.

## Financial support and sponsorship

Nil.

## Conflicts of interest

There are no conflict of interests.

---

## References

- 1 Gerfard-Valentin M, Maucort-Boulch D, Hot A, Iwas J, Ninet J, Durieu I *et al*. Adult onset Still's disease: manifestations, treatments, outcome and prognostic factors in 57 patients. *Medicine (Baltimore)* 2014; 93:91–99.
- 2 Magadur-Joly G, Billaud E, Barrier JH, Pennec YL, Masson C, Renou P *et al*. Epidemiology of adult onset Still's disease: estimate of the incidence by a retrospective study in west France. *Ann Rheum Dis* 1995; 54:587–590.
- 3 Yamaguchi M, Ohta A, Tsunematsu T, Kasukawa R, Mizushima Y, Kashiwagi H, *et al*. Preliminary criteria for classification of adult Still's disease. *J Rheumatol* 1992; 19:424–430.
- 4 Ohashi M, Moriya C, Kanoh H, Kinomura Y, Terakura Y, Kasahara S, *et al*. Adult-onset Still's disease with dermatomyositis-like eruption. *Dermatology* 2012; 39:958–960.
- 5 Uppal SS, Al-Mutairi M, Hayat S, Abraham M, Malaviya A. Ten years of clinical experience with adult onset Still's disease: is the outcome improving? *Clin Rheumatol* 2007; 26:1055–1060.
- 6 Jamilloux Y, Gerfard-Valentin M, Henry T, Sève P. Treatment of adult-onset Still's disease: a review. *Ther Clin Risk Manag* 2015; 11:33–43.
- 7 Zong M, Dorph C, Dastmalchi M, Alexanderson H, Pieper J, Amoudruz P *et al*. Anakinra treatment in patients with refractory inflammatory

- myopathies and possible predictive response biomarkers: a mechanistic study with 12 months follow-up. *Ann Rheum Dis* 2014; 73:913–920.
- 8 Jie W, Miao L, Yankun S, Hong Y, Zhongxin X. Demyelinating encephalopathy in adult onset Still's disease: case report and review of literature. *Clin Neurol Neurosurg* 2013; 115:2213–2216.
  - 9 Desai SS, Allen E, Deodhar A. Miller Fisher syndrome in adult onset Still's disease: case report and review of literature of other neurological manifestations. *Rheumatology* 2002; 41:216–222.
  - 10 Dabadghao VS, Sharma SK, Mangudkar SS, Garg SA. Seizures in a patient of adult onset Still's disease. *Med J DY Patil Univ* 2014; 7:400–402.